



Essential Skills Evaluation: Pretest and Posttest from pages 4–20

- 1) $\frac{1}{2}$; 3 times a day 2) 3; 2 times a day 3) 10; 2; once a day 4) $\frac{1}{2}$; every 3 hours as needed for moderate pain
 5) 2; once every morning 6) 2.5; every 8 hours




© Cengage Learning 2013

- 7) 0.5; every 4 hours as needed for nausea




© Cengage Learning 2013

- 8) 0.8; once, immediately




© Cengage Learning 2013

- 9) 0.4; every 4 hours as needed for severe pain



© Cengage Learning 2013


- 10) 1.5; once, immediately



© Cengage Learning 2013


Essential Skills Evaluation — Pages 4–20

- 11) 1.4; every 8 hours




© Cengage Learning 2013

- 12) 0.88; once, immediately




© Cengage Learning 2013

- 13) 1; once, immediately; either syringe is appropriate




© Cengage Learning 2013




© Cengage Learning 2013

- 14) 0.5; once, every morning; either syringe is appropriate




© Cengage Learning 2013




© Cengage Learning 2013

15) 8; every 12 hours




© Cengage Learning 2013

16) 4; every 12 hours




© Cengage Learning 2013

17) 1; 1; 0.25; every 8 hours




© Cengage Learning 2013

18) 1.4; 75; every 6 hours



© Cengage Learning 2013


- 19) 68; once a day before breakfast




© Cengage Learning 2013

- 20) 1

- 21) Benadryl; 0.7; either syringe is appropriate




© Cengage Learning 2013



© Cengage Learning 2013


- 22) Narcan; 1



© Cengage Learning 2013

- 23) Yes. Her temperature is 102.2°F. Tylenol every 4 hours as needed is indicated for fever greater than 101°F. It has been 5 hours and 5 minutes since her last dose. 24) 2 25) 18.8; 138 26) 138 27) 8; 2 28) 1; 0700; 1200; 1700

- 29) 18; subcutaneous



© Cengage Learning 2013

- 30) 113

- 31) Yes; the usual dosage is 20–40 mg/kg/day divided into 3 doses q.8h, which is equivalent to 67–133 mg per dose for a 22 lb (10 kg) child.